

# CHOOSE CONFIDENCE

## With *Natrelle*<sup>®</sup> Cohesive Gel



Actual *Natrelle*<sup>®</sup> Gel Style 10 Breast Implant Cut in Half\*

- Only *Natrelle*<sup>®</sup> Gel implants are guaranteed for life and automatically enrolled in the *ConfidencePlus*<sup>®</sup> Premier Warranty program<sup>1</sup>
- *Natrelle*<sup>®</sup> uses the latest generation of FDA-approved cohesive silicone gel<sup>2</sup>

**Visit [Natrelle.com](http://Natrelle.com) for more information**

\*Implant may react differently if shell is breached while implanted

<sup>1</sup> The *Natrelle*<sup>®</sup> *ConfidencePlus*<sup>®</sup> Premier Warranty includes free product replacement (including contralateral implants) for implant deflation or rupture due to loss of shell integrity for the lifetime of the device. Also includes surgery reimbursement up to \$3,500 for up to 10 years. Visit <http://www.cppwarranty.com> for complete coverage details.

<sup>2</sup> <http://www.breastimplantanswers.com>



Scan this box for more information.

### Important *Natrelle*<sup>®</sup> Safety Information

The *Natrelle*<sup>®</sup> Collection of Breast Implants is indicated for females for breast augmentation and breast reconstruction. Breast Augmentation includes primary breast augmentation to increase the breast size, as well as revision surgery to correct or improve the result of a breast augmentation surgery. Breast Reconstruction includes primary reconstruction to replace breast tissue that has been removed due to cancer or trauma or that has failed to develop properly due to severe abnormality as well as revision surgery to correct or improve the result of a breast reconstruction surgery. For breast augmentation, *Natrelle*<sup>®</sup> silicone filled breast implants are indicated for women at least 22 years old and *Natrelle*<sup>®</sup> saline filled breast implants are indicated for women at least 18 years old.

Complications may include, among others, deflation or rupture, reoperation, capsular contracture, breast pain, and swelling. Contraindicated in women with active infection, pre or existing breast cancer who have not received adequate treatment, and in women who are pregnant or nursing. For more detailed information now, please read the complete patient information available at [http://natrelle.com/safety\\_information](http://natrelle.com/safety_information).

